



# Bethlem myopathy

## What is Bethlem myopathy?

Bethlem myopathy falls under the category of congenital muscular dystrophies, within a sub-type known as the 'collagen VI-related myopathies'. Bethlem myopathy has the following specific features:

- the joints of the hands develop 'contractures' or 'tightness' at the level of the fingers, and the feet develop 'contractures' or 'tightness' of the Achilles tendons
- a variable degree of muscle weakness, affecting the legs especially and to a lesser extent the arms, develops and slowly progresses over time
- respiratory muscles are more mildly affected, and problems related to respiratory insufficiency are very rare and limited to the late adult years.

## What causes Bethlem myopathy?

Bethlem myopathy is caused by mutations in at least three genes: *COL6A1*, *COL6A2* and *COL6A3*. These genes are responsible for the production of a protein called collagen VI, which is required to maintain the structure of the muscles.

## How is Bethlem myopathy diagnosed?

The diagnosis of Bethlem myopathy can usually be suspected from an individual's medical history and examination. The specific diagnosis is generally made by looking at a piece of muscle or sometimes skin; this is called a biopsy. However, before doing a muscle biopsy, a few other tests may be done.

A **blood test** can measure the level of a muscle protein (called creatine kinase or CK). In people with Bethlem myopathy, the CK level is either normal or mildly elevated.

A **muscle biopsy** involves taking out a small piece of muscle, usually from the thigh. The piece of muscle is then studied under the microscope, looking for signs which might indicate the type of abnormality that characterises the condition. It is also possible to look at the expression of collagen VI in the muscle under the microscope. There are specific 'markers' or 'tags' which can detect if collagen VI is normally present or reduced.

**Genetic tests** looking for abnormalities in one of the three genes responsible for Bethlem myopathy (*COL6A1*, *COL6A2* and *COL6A3*) are now available in the UK. These are done in a specific, nationally-designated central laboratory (NSCT Centre for CMD and congenital myopathies, London) and can provide a definitive diagnosis.

This factsheet is under review, due for updating later in 2017. If you have any queries, please contact us.

Version 3 / Date published: March 2003 / Original author: Dr Reghan Foley, Dr Stephanie Robb and Prof Francesco Muntoni, Dubowitz Neuromuscular Centre  
Updated: October 2013 / Updated by: Dr Reghan Foley / Date of review: October 2014  
Registered Charity No. 205395 and Registered Scottish Charity No. SC039445





## Is there a treatment or cure?

At present there is no cure for Bethlem myopathy but there are ways, described below, of helping to alleviate the effects of the condition. Research into the congenital muscular dystrophies is nevertheless developing, and it is likely that clinical trials will start in the not-too-distant future.

Physiotherapy is one of the main forms of help. An initial physiotherapy assessment at the time of diagnosis should be followed by an exercise programme and regular check-ups. The main aim of physiotherapy is to keep the muscles as active as possible and to prevent or slow the progression of joint contractures (tightness). Individuals with Bethlem myopathy are encouraged to remain as active as possible. Swimming is a particularly good form of exercise. Physiotherapists can also provide advice on orthoses, walking aids and a wheelchair, when necessary.

Occasionally, surgery to release the Achilles tendons can help individuals with Bethlem myopathy to stand and walk more easily.

Children and adults with Bethlem myopathy should ideally regularly visit a specialist neuromuscular clinic, with access to physiotherapy, orthotic, respiratory, orthopaedic and genetic specialists as needed.

## What is the prognosis?

The first symptoms of Bethlem myopathy can present at any time from birth through to adulthood and are variable. In childhood, these symptoms can be hypotonia (floppiness) and joint laxity (double jointedness), muscle weakness, delayed achievement of motor milestones such as sitting unaided or walking. Joint contractures are common in Bethlem myopathy, typically affecting the wrists, fingers, elbows and Achilles tendons. Some stiffness of the spine can also develop over the years.

Other symptoms such as poor stamina/exercise intolerance and difficulty rising from a seated position and climbing stairs are related to the proximal muscle weakness evident in Bethlem myopathy.

As collagen VI is found in the skin as well as the muscles, individuals with Bethlem myopathy have a tendency for scars which heal slowly or become thickened and elevated (keloid formation). The surfaces of the arms and legs can feel 'rough' or 'bumpy' owing to 'keratosis pilaris' which commonly occurs in the skin.

The main complications in Bethlem myopathy are the progression of joint contractures (tightness), increased proximal limb (hip and shoulder girdle) weakness and, to a much lesser extent, a decline in lung function. Physiotherapy and a regimen of stretching can help to prevent or slow the progression of joint contractures. Some adults will become dependent on walking aids (cane, crutches or wheelchair) for walking outdoors.

Lung function in individuals with Bethlem myopathy should be monitored. If indicated (depending on the respiratory function values), overnight sleep studies could be required.

This factsheet is under review, due for updating later in 2017. If you have any queries, please contact us.

Version 3 / Date published: March 2003 / Original author: Dr Reghan Foley, Dr Stephanie Robb and Prof Francesco Muntoni, Dubowitz Neuromuscular Centre  
Updated: October 2013 / Updated by: Dr Reghan Foley / Date of review: October 2014  
Registered Charity No. 205395 and Registered Scottish Charity No. SC039445



Rarely, adults with Bethlem myopathy develop sleep hypoventilation for which night-time non-invasive ventilation (NIV) will be necessary.

## Other related publications

- Congenital muscular dystrophies
- MDC1A (merosin-deficient congenital muscular dystrophy)
- SEPN1-related myopathy
- Ullrich congenital muscular dystrophy
- Carrier detection tests and prenatal diagnosis of inherited neuromuscular conditions
- Inheritance and the muscular dystrophies
- Muscle biopsies
- Surgical correction of spinal deformity in muscular dystrophy and other neuromuscular disorders

## We're here for you at the point of diagnosis and at every stage thereafter, and can:

- ▶ give you accurate and up-to-date information about your or your child's muscle-wasting condition, and let you know of progress in research
- ▶ give you tips and advice about day-to-day life, written by people who know exactly what it's like to live with a muscle-wasting condition
- ▶ put you in touch with other families living with the same muscle-wasting condition, who can tell you about their experiences
- ▶ tell you about – and help you get – the services, equipment and support you're entitled to.

If you have feedback about this factsheet please email [info@muscular dystrophyuk.org](mailto:info@muscular dystrophyuk.org).

### Disclaimer

While every reasonable effort is made to ensure that the information in this document is complete, correct and up-to-date, this cannot be guaranteed and Muscular Dystrophy UK shall not be liable whatsoever for any damages incurred as a result of its use. Muscular Dystrophy UK does not necessarily endorse the services provided by the organisations listed in our factsheets.

## Here for you

The friendly staff in the care and support team at the Muscular Dystrophy UK's London office are available on **0800 652 6352** or [info@muscular dystrophyuk.org](mailto:info@muscular dystrophyuk.org) from 8.30am to 6pm Monday to Friday to offer free information and emotional support.

If they can't help you, they are more than happy to signpost you to specialist services close to you, or to other people who can help.

**[www.muscular dystrophyuk.org](http://www.muscular dystrophyuk.org)**

This factsheet is under review, due for updating later in 2017. If you have any queries, please contact us.

Version 3 / Date published: March 2003 / Original author: Dr Reghan Foley, Dr Stephanie Robb and Prof Francesco Muntoni, Dubowitz Neuromuscular Centre  
Updated: October 2013 / Updated by: Dr Reghan Foley / Date of review: October 2014  
Registered Charity No. 205395 and Registered Scottish Charity No. SC039445

